

## Anti-RNF36/TRIM69 antibody (C-Term) (STJ70340) STJ70340

## **GENERAL INFORMATION**

Host/Source Goat

Product Type Primary antibodies Short Goat polyclonal antibody anti-RNF36/TRIM69 (C-Term) is suitable for use in ELISA, Western Blot and Immunohistochemistry **Description** research applications. Applications Pep-ELISA/WB/IHC Reactivity Human/Mouse/Rat/Dog/Pig/Cow

## **PRODUCT PROPERTIES**

| Clonality<br>Clone ID | Polyclonal   |
|-----------------------|--|
| Concentration         | 0.5 mg/mL  |
| Conjugation           | Unconjugated   |
| Purification          | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing |
|                       | peptide.   |
| Dilution Range        | Peptide ELISA: antibody detection limit dilution 1:32000.  |
|                       | WB: Approx 55kDa band observed in Mouse Testis lysates (calculated MW of 57.3kDa according to Mouse NP_536771.2).            |
|                       | Recommended concentration: 0.5-2µg/ml.   |
|                       | IHC: Paraffin embedded Human Te  |
| Formulation           | 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA                                       |
| Isotype               | IgG  |
| Storage               | Store at-20°C on receipt and minimise freeze-thaw cycles.  |
| Instruction           |  |

## **TARGET INFORMATION**

| Gene ID     |   |
|-------------|---|
| Gene Symbol |   |
| Uniprot ID  | TRI69_HUMAN   |
| Immunogen   |   |
| Immunogen   | C-Term  |
| Region      |   |
| Specificity | This antibody is expected to recognize both reported isoforms (NP_892030.3; NP_542783.2). |
|             | GENKEPLHILHPQ   |
| Sequence    |   |
|             |   |
| 250kDa      |   |
| 150kDa      |   |
| 100kDa      |   |
| 75kDa       |   |
| 50kDa       |   |
| 37kDa       |   |
|             |   |
| 25kDa       |   |
| ZokDa       |   |
| 20kDa       |   |
|             |   |

15kDa STJ70340 (0. 5ŵg/m) staining of Mouse Testis lysate (35ŵg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081