

## Anti-AIP/ARA9 antibody (C-Term) (STJ71174) STJ71174

## **GENERAL INFORMATION**

Product Type	Ρ
Short	G
Description	а
Applications	Ρ
Host/Source	G
Reactivity	Н

Primary antibodies Goat polyclonal antibody anti-AIP/ARA9 (C-Term) is suitable for use in ELISA, Western Blot and Immunohistochemistry research applications. Pep-ELISA/WB/IHC Goat

Reactivity Human/Mouse/Rat/Dog

## **PRODUCT PROPERTIES**

Clonality Clone ID	Polyclonal
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing
	peptide.
Dilution	Peptide ELISA: antibody detection limit dilution 1:64000.
Range	WB: Approx 37kDa band observed in lysates of cell line NIH/3T3 (calculated MW of 37.7kDa according to Human NP_003968.1 and
	37.6kDa according to mouse NP_057875.1). In transfected HEK293 t
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage	Store at-20°C on receipt and minimise freeze-thaw cycles.
Instruction	

## **TARGET INFORMATION**

STJ711 (35µg 1 hour.

Gene S Unij Immu Immu Spe Immu	inogen inogen Region cificity	AIP AIP_HUMAN	DKAR			
	250kDa 150kDa 100kDa 75kDa		188 98			
	50kDa 37kDa		62	-		
			49 38	_	-	
	25kDa 20kDa		28 17 14			
174 (0. 02µg/ml) protein in RIPA b Detected by chen	uffer). Primary	incubation was	HEK293 overexpress with STJ71174 (mocl by Origene.	ing ARA9		

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081