

## Anti-BOC antibody (Internal) (STJ71520)

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Goat polyclonal antibody anti-BOC (Internal) is suitable for use in ELISA and Western Blot research applications.

Description

Applications Pep-ELISA/WB

Host/Source Goat

Reactivity Human/Mouse/Dog/Cow

## **PRODUCT PROPERTIES**

Clonality Polyclonal Clone ID

Concentration 0.5 mg/mL Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

**Dilution Range** Peptide ELISA: antibody detection limit dilution 1:2000.

WB: Approx 110kDa band observed in Human Heart and Ileum lysates (calculated MW of 121kDa according to NP\_150279.1).

Recommended concentration: 0.3-1µg/ml.

Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

## **TARGET INFORMATION**

Gene ID 91653 Gene Symbol BOC

Uniprot ID BOC\_HUMAN

Immunogen

Immunogen Internal Region

Specificity
Immunogen TQEVKTSGSSDRLR

Sequence

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

STJ71520 (0. 3µg/ml) staining of Human Ileum lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.