

Anti-TAP1 antibody (Internal) (STJ71702) STJ71702

GENERAL INFORMATION


Host/Source Goat Reactivity Human

Product Type Primary antibodies Short Description Goat polyclonal antibody anti-TAP1 (Internal) is suitable for use in ELISA and Western Blot research applications. Applications Pep-ELISA/WB

PRODUCT PROPERTIES

Clonality Polyclonal Clone ID Concentration 0.5 mg/mL Conjugation Unconjugated Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Dilution Range Peptide ELISA: antibody detection limit dilution 1:32000. WB: Approx 85kDa band observed in lysates of cell line MCF7 (calculated MW of 87.2kDa according to NP_000584.2). Recommended concentration: 1-3µg/ml. Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA Isotype IgG Storage Store at-20°C on receipt and minimise freeze-thaw cycles. Instruction

TARGET INFORMATION



15kDa STJ71702 (1µg/ml) staining of MCF7 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081