

Anti-OXT-Mouse antibody (C-Term) (STJ72156)

STJ72156

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-OXT-Mouse (C-Term) is suitable for use in ELISA, Immunofluorescence and Flow Cytometry research

 Description
 applications.

 Applications
 Pep-ELISA/IF/FC

 Host/Source
 Goat

 Reactivity
 Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration 0.5 mg/mL

Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

Dilution Range Peptide ELISA: antibody detection limit dilution 1:16000.

WB: Preliminary experiments gave an approx 22kDa band in Mouse Liver after 1µg/ml antibody staining. Please note that currently

we cannot find an explanation in the literature for this ban

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

TARGET INFORMATION


Gene ID Gene Symbol Uniprot ID Immunogen

Immunogen C-Term


Region Specificity

Immunogen DGCRTDPACDPES

Sequence



STJ/2156 Immunofluorescence analysis o paraformalchyde fixed NIH3T3 cells, permeabilizer with 0. 15% Triton, Primary incubation Thr (10ug/ml followed by Alexa Fluor 488 secondary antibod-(2ug/ml), showing cytoplasmic and membrane staining The nuclear stain is DAPI (blue). Negative control Unimmunized goat IgG (10ug/ml) followed by Alexi Fluor 488 secondary antibod-vizu/ml).



STJ72156 Flow cytometric analysis o paraformaldehyde fixed NIH3T3 cells (blue line) permeabilized with 0.5% friton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG contol: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.