

Anti-NEK7 antibody (Internal) (STJ72425) STJ72425

GENERAL INFORMATION

••	Primary antibodies Goat polyclonal antibody anti-NEK7 (Internal) is suitable for use in ELISA, Western Blot and Immunohistochemistry research
Description	applications.
Applications	Pep-ELISA/WB/IHC
Host/Source	Goat
Reactivity	Human/Mouse/Rat/Dog/Cow

PRODUCT PROPERTIES

Clonality Clone ID	Polycional			
Concentration	0.5 mg/mL			
Conjugation	Unconjugated			
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.			
Dilution	Peptide ELISA: antibody detection limit dilution 1:64000.			
Range	WB: Approx 35kDa band observed in lysates of cell line HeLa (calculated MW of 34.6kDa according to NP_598001.1). Recommended concentration: 0.05-0.2µg/ml. Primary incubation was 1 hour.			
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA			
Isotype	lgG			
Storage Instruction	Store at-20°C on receipt and minimise freeze-thaw cycles.			

TARGET INFORMATION

Gene Sy Unip Immui Immui R Spec Immui	nogen nogen legion cificity	
1 1 7 5	250kDa 50kDa 00kDa 55kDa 50kDa 87kDa	
2	25kDa 20kDa	

15kDa STJ72425 (0, 05ŵg/m) staining of HeLa lysate (35ŵg protein in RIPA buffer). Detected by chemiluminescence.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081