

Anti-PRDM14 antibody (414-427) (STJ72514)

STJ72514

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-PRDM14 (414-427) is suitable for use in ELISA, Immunofluorescence and Flow Cytometry research

Description applications.

Applications Pep-ELISA/IF/FC
Host/Source Goat

Reactivity Human/Mouse/Rat/Dog/Pig/Cow

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

ID

Concentration 0.5 mg/mL **Conjugation** Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

Dilution Peptide ELISA: antibody detection limit dilution 1:16000.

Range WB: Preliminary experiments gave an approx 40kDa band in Human Heart lysates after 0.1-1µg/ml antibody staining and this band

was successfully blocked by incubation with the immunizing pep

Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction


TARGET INFORMATION

Gene ID 63978
Gene Symbol PRDM14
Uniprot ID PRD14_HUMAN


Immunogen Immunogen 414-427 Region

Specificity Immunogen YRDKHLKYTPCVDK

Sequence



STJ72514 Immunofluorescence analysis or paraformatidelytide fixed He.d. costle, permeabilized with 0. 15% Triton. Pirmay incubation The (Tolughi followed by Alex Fliuor 488 secondary antibod (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain in DAPI (blue). Negative control: Unimmunized goat [of (Toughin) followed by Alexa Fliuor 488 secondar



STJ72514 Flow cytometric analysis o paraformaldehyde fixed HeLa cells (blue line) permeabilized with 0.5% firton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondar antibody (1ug/ml), IgG control: Unimmunized goal IgG (black line) followed by Alexa Fluor 488 secondary antibody.