

## Anti-Uromodulin antibody (525-536) (STJ72846)

STJ72846

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Goat polyclonal antibody anti-Uromodulin (525-536) is suitable for use in ELISA and Immunofluorescence research applications.
<b>Applications</b>	Pep-ELISA/IF
<b>Host/Source</b>	Goat
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	0.5 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:64000. IF: Strong expression of the protein seen in the Plasma membrane and Golgi of HeLa cells. Recommended concentration: 10µg/ml.
<b>Formulation</b>	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C on receipt and minimise freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	7369
<b>Gene Symbol</b>	UMOD
<b>Uniprot ID</b>	UROM_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	525-536
<b>Specificity</b>	Reported variants represent identical protein: NP_001008390.1, NP_003352.2
<b>Immunogen Sequence</b>	DRCPHTRDSTIQ



STJ72846 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml), showing membrane and Golgi staining. The nuclear stain is DAPI (blue). NA NA Negative control: Unimmunized goat IgG (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml).

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081