

## Anti-periostin antibody (C-Term) (STJ72902)

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-periostin (C-Term) is suitable for use in ELISA and Western Blot research applications.

Applications Pep-ELISA/WB Host/Source Goat Reactivity Human

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

Concentration 0.5 mg/mL Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

Peptide ELISA: antibody detection limit dilution 1:16000. WB: Approx 100kDa band observed in lysates of cell line MCF7 (calculated MW of 93.3kDa according to NP\_006466.2).

Recommended concentration: 1-3µg/ml.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$ 

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

**Dilution Range** 

## **TARGET INFORMATION**

Gene ID 10631

Gene Symbol POSTN

Uniprot ID POSTN\_HUMAN

Immunogen

Immunogen C-Term Region

Specificity This antibody is expected to recognize all reported isoforms (NP\_006466.2; NP\_001129406.1; NP\_001129407.1;

NP\_001129408.1).

Immunogen RKLQANKKVQGSRR

Sequence

250kDa 150kDa

100kDa

75kDa 50kDa

37kDa

25kDa 20kDa

15kDa STJ72902 (2Åug/ml) staining of MCF7 lysate (35Åug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.