

Anti-NDRG1, Biotinylated antibody (C-Term) {Biotin} (STJ73261)

STJ73261

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-NDRG1, Biotinylated (C-Term) is suitable for use in ELISA and Western Blot research applications.

Description

Applications Pep-ELISA/WB Host/Source Goat Reactivity Human

PRODUCT PROPERTIES

Clonality Polyclonal
Clone ID

Concentration 0.5 mg/mL
Conjugation Biotin

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

Dilution Peptide ELISA: antibody detection limit dilution 1:32000.

Range WB: Approx 48kDa band observed in Human Kidney lysates (calculated MW of 42.8kDa according to NP_006087.2). See non-

biotinylated parental product's datasheet for further QC data. Recommend

Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 10397 Gene Symbol NDRG1

Uniprot ID NDRG1_HUMAN

Immunogen Immunogen C-Term

Region C-Ten

Region

Specificity Reported variants represent identical protein: NP_006087.2, NP_001128714.1

Immunogen GNSAGPKSMEVSC.

Sequence

250kDa 150kDa 100kDa

> 50kDa 37kDa

> > 25kDa 20kDa

Biotinylated STJ70275 (ZApy/ml) staining of Human Kidney Iysate (35Apy protein in RIPA buffer) , exactly mirroring its parental non-biotinylated product. Primar incubation was I hour. Detected by homelly being career, using street wide, MIDP and using the production of the control of t

chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.