

Anti-MYH9 antibody (Internal) (STJ73316) STJ73316

GENERAL INFORMATION

Product Type Primary antibodies Description Applications Pep-ELISA/WB Host/Source Goat

Short Goat polyclonal antibody anti-MYH9 (Internal) is suitable for use in ELISA and Western Blot research applications. Reactivity Human/Mouse/Rat/Dog

PRODUCT PROPERTIES

Clonality Clone ID	Polyclonal
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Dilution	Peptide ELISA: antibody detection limit dilution 1:32000.
Range	WB: Approx 250kDa band observed in lysates of Human Peripheral Blood Mononucleocytes and of NIH3T3 (calculated MW of 227kDa
	according to Human NP_002464.1 and to Mouse NP_071855.2). Recomm
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage	Store at-20°C on receipt and minimise freeze-thaw cycles.
Instruction	

TARGET INFORMATION

15kDa

STJ73316 (0. 3µg/ml) staining blood lymphocytes lysate (35Â buffer). Primary incubation was

Human peripheral protein in RIPA

Immunogen Immunogen Region Specificity	MYH9 MYH9_HUMAN	
 250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 		250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa
15kDa		15kDa

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081

STJ73316 (1µg/ml) staining of NIH3T3 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.