

## Anti-HMGN2 antibody (1-50 aa) (STJ93550)

STJ93550

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Non-histone chromosomal protein HMG-17 (1-50 aa) is suitable for use in Immunohistochemistry, Immunofluorescence and ELISA research applications.
<b>Applications</b>	IHC/IF/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution Range</b>	IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:40000
<b>Formulation</b>	Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	3151
<b>Gene Symbol</b>	HMGN2
<b>Uniprot ID</b>	HMGN2_HUMAN
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the human HMG17 at the amino acid range 1-50
<b>Immunogen</b>	1-50 aa
<b>Region</b>	
<b>Specificity</b>	HMG-17 Polyclonal Antibody detects endogenous levels of HMG-17 protein.
<b>Immunogen Sequence</b>	



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using HMG17 Antibody. The picture on the right is blocked with the synthesized peptide.

Immunofluorescence analysis of HeLa cells, using HMG17 Antibody. The picture on the right is blocked with the synthesized peptide.