

Anti-HNRNPK antibody (250-299 aa) (STJ93567) STJ93567

GENERAL INFORMATION

 Product Type
 Primary antibodies

 Short
 Rabbit polyclonal antibody anti-Heterogeneous nuclear ribonucleoprotein K (250-299 aa) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and ELISA research applications.

 Applications
 WB/IHC/IF/ELISA

 Reactivity
 Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Clone ID	Polyclonal				
Concentration	1 mg/mL				
Conjugation	Unconjugated				
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.				
Dilution	WB 1:500-1:2000				
Range	IHC 1:100-1:300				
	ELISA 1:40000				
	IF 1:50-200				
Formulation	Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.				
Isotype	IgG				
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.				

TARGET INFORMATION

TARGET INFORMATION					
Gene ID Gene Symbol Uniprot ID Immunogen Region Specificity Immunogen Sequence	HNRNPK HNRPK_HUN The antiserur 250-299 aa		e derived from the human hnRNP K at the ar	nino acid range 250-299	
K562 A549HT29 A5	549			HepG2	
	117		and the second	(kD)	
	85		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	117- 85-	
hnRNPK	48		A & P & P & P & P & P & P & P & P & P &	48-	
	34	a geographic states	1	34-	
	26	and the second second	Contraction of the second second	26-	
	19			19-	
	(kD)	Immunohistochemical analysis of paraffin-embedded	Immunchistochemical analysis of paraffin-embedded		
Western blot analysis of lysates from K562, A549, and HT-29 cells, using hnRNP K Antibody. The lane on the right is blocked with the synthesized peptide.		Immunchistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4ŰC overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negetive contri (right) obtaned from antibody was pre-absorbed by immunogen peptide.	Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4ŰC overnight). High-pressure and temperature Tris-EDTA, pH8.0 was used for antigen retrieval. Negetive contri (right) obtaned from antibody was pre-absorbed by immunogen peptide.	Western blot analysis of various cells using hnRNP K Polyclonal Antibody diluted at 111/4 2000	

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081