

Anti-ANGPTL4 antibody (301-350 aa) (STJ96539)

STJ96539

GENERAL INFORMATION

Product Type Primary antibodies

Short Rabbit polyclonal antibody anti-Angiopoietin-related protein 4 (301-350 aa) is suitable for use in Western Blot,

Description Immunohistochemistry, Immunofluorescence and ELISA research applications.

Applications WB/IHC/IF/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration 1 mg/mL

Conjugation Unconjugated

Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Dilution Range IHC-P 100-300 WB 1:500-1:2000

ELISA 1:20000

IF 1:50-200

Formulation Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 51129 ene Symbol ANGPT

Gene Symbol ANGPTL4

Uniprot ID ANGL4_HUMAN

Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human ANGPTL4 at the amino acid


range 301-350 **Immunogen** 301-350 aa

Region


Specificity Immunogen

Angptl4 Polyclonal Antibody detects endogenous levels of Angptl4 protein.

Immunogen Sequence



Western blot analysis of lysate from MCF7 cells, using



Human testis. 1. Antibody was diluted at 1:100 (4Å*C pernight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

Himman testis. 1. Antibody was diluted at 1:100 (4A°C overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

nmunonistochemical analysis of paramin-embedged uman testis. 1, Antibody was diluted at 1:100 (4A°C vernight). 2, High-pressure and temperature EDTA, H8.0 was used for antigen retrieval. 3, Secondary titibody was diluted at 1:200 (room temperature, Omin).