

Anti-CXCL13 antibody (41-90 aa) (STJ96573)

STJ96573

GENERAL INFORMATION

Product Type Primary antibodies

Short Rabbit polyclonal antibody anti-C-X-C motif chemokine 13 (41-90 aa) is suitable for use in Western Blot, Immunohistochemistry,

Description Immunofluorescence and ELISA research applications.

Applications WB/IHC/IF/ELISA

Host/Source Rabbit

Reactivity Human/Rat/Mouse

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration 1 mg/mL

Conjugation Unconjugated

Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Dilution Range WB 1:500-1:2000

IHC-P 1:100-300 ELISA 1:20000

IF 1:50-200

Formulation Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 10563

Gene Symbol CXCL13

Uniprot ID CXL13_HUMAN

Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human CXCL13 at the amino acid


range 41-90

Immunogen 41-90 aa


Region

Specificity BLC Polyclonal Antibody detects endogenous levels of BLC protein. **Immunogen**

Sequence



Western blot analysis of lysate from mouse spleen cells, using CXCL13 Antibody.



Human Liver. 1, Antibody was diluted at 1:200 (4ŰC overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

Human Liver. 1, Antibody was diluted at 1:200 (AAYC overnight). 2, High-pressure and temperature EDTA pH8.0 was used for antigen retrieval. 3, Secondar antibody was diluted at 1:200 (room temperature 30min).

Human Liver. 1, Antibody was diluted at 1:200 (AA'C online). 2, High-pressure and temperature EDTA pH8.0 was used for antigen retrieval. 3, Secondar antibody was diluted at 1:200 (room temperature 30min).