

Anti-Nestin antibody [SM4923] (STJA0004923)

STJA0004923

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Mouse monoclonal antibody anti-Nestin is suitable for use in Immunohistochemistry and Immunofluorescence research

applications.

Applications IHC/IF Host/Source Mouse

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clone ID Monoclonal SM4923

Concentration

Conjugation Unconjugated Purification Affinity purification

Dilution Range IHC/IF (H, M, R) 1:500-1:1000/1:500-1:1000

Formulation PBS with 0.02% sodium azide, 100 Mu g/ml BSA and 50% glycerol.

Isotype IgG

Storage Instruction Store at-20C for up to one year, and avoid repeated freeze-thaw cycles.

TARGET INFORMATION


Gene ID 18008 Gene Symbol Nes

Uniprot ID NEST_MOUSE


Immunogen Immunogen Region Specificity Immunogen

Sequence


Immunogen Recombinant protein corresponding to Mouse Nestin




Immunohistochemistry analysis of Nestin. Sampl Human kidney, 4% PFA 12-24h, Antigen retrieva Citrate buffer, pressure cooker 2min. Blocking buffe 3% BSA in PBS, RT, 30min. Primary antibody: 1:500 4C overnight. Secondary antibody: HRP Goat Ant



mmunohistochemistry analysis of Nestin Samplemouse kidney, 4% PFA 1-2-4h, Antigen retireal Citrate buffer, pressure cooker 2min, Blocking buffer 3% BSA in PBS, RT, 30min, Primary antibody. 1:500 to overnight. Secondary antibody: HRP Goat Ant



Immunohistochemistry analysis of Nestin. Sample: Ridney. 4% PFA 12-24h. Antigen retrieval: Citra buffer, pressure cooker 2min. Blocking buffer; 3% BS in PBS, FTT, 30min. Primary antibody: 1:500, 4 overnight. Secondary antibody: HRP Goat Anti-mou



IF analysis of Nestin. Sample: mouse kidney, 4% PFA 12-24h. Antigen retrieval: Citrate buffer, pressure cooker Zmin. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1:500, 4C overnight. Secondary antibody: Cy3 conjugated Goat Anti-mouse IgG, 1:200 RT 1h.