

## Goat Anti-Human IgG Fab Fragment antibody {Biotin} (STJS000146)

STJS000146

### GENERAL INFORMATION

|                          |  |
|--------------------------|--|
| <b>Product Type</b>      | Secondary antibodies   |
| <b>Short Description</b> | Biotin-conjugated goat polyclonal anti-Human IgG Fab Fragment secondary antibody. For use in most research applications. |
| <b>Applications</b>      | ELISA/WB/IHC/ICC   |
| <b>Host/Source</b>       | Goat   |
| <b>Reactivity</b>        | Human  |

### PRODUCT PROPERTIES

|                            |  |
|----------------------------|--|
| <b>Clonality</b>           | Polyclonal   |
| <b>Clone ID</b>            |  |
| <b>Concentration</b>       | 1 mg/mL  |
| <b>Conjugation</b>         | Biotin   |
| <b>Purification</b>        | The antibody was isolated from ascitic by immunoaffinity chromatography using antigens coupled to agarose beads. |
| <b>Dilution Range</b>      | ELISA 1:5000-1:20000<br>WB 1:5000-1:20000<br>IHC 1:500-1:1000<br>ICC 1:500-1:1000                                |
| <b>Formulation</b>         | Liquid in 0.01M PBS pH7.2, 1% BSA, 50% Glycerol and 0.05% Sodium Azide   |
| <b>Isotype</b>             | IgG  |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                   |

### TARGET INFORMATION

|                         |
|-------------------------|
| <b>Gene ID</b>          |
| <b>Gene Symbol</b>      |
| <b>Uniprot ID</b>       |
| <b>Immunogen</b>        |
| <b>Immunogen Region</b> |
| <b>Specificity</b>      |
| <b>Immunogen</b>        |
| <b>Sequence</b>         |