

Goat Anti-Rabbit IgG H+L antibody {Cy3} (STJS001166)

STJS001166

GENERAL INFORMATION


Product Type	Secondary antibodies
Short Description	
Applications	IF/ICC/FC
Host/Source	Goat
Reactivity	Rabbit

PRODUCT PROPERTIES



Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Cy3
Purification	Affinity purification
Dilution Range	IF/ICC:1:100-1:800 FC:1:100-1:800
Formulation	PBS with 0.025% Sodium Azide, 0.75% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	

TARGET INFORMATION

Gene ID	
Gene Symbol	
Uniprot ID	
Immunogen	
Immunogen Region	
Specificity	Rabbit IgG
Immunogen Sequence	



Confocal imaging of U-2 OS cells using DNA topoisomerase I (TOP1) Rabbit mAb (STJ235679, dilution 1:100) (Red). The cells were counterstained with Alpha-Tubulin Mouse mAb (STJ11107558, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (STJS001166, dilution 1:100) (Red), ABTolA® 488-conjugated Goat Anti-Mouse IgG (H+L) (STJS001209, dilution 1:200) (Green)



Flow cytometry: 1X10^6 K-562 cells (negative control, left) and A-431 cells (right) were surface-stained with Purified Rabbit anti-Human E-Cadherin mAb (5 Mu l/Test, orange line) or secondary antibody only (blue line). Non-fluorescently stained K-562 and A-431 cells were used as blank control (red line). Cy3 Goat Anti-Rabbit IgG (H+L) (STJS001166, 1:800) was used as a secondary antibody.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081