

Goat Anti-IgG H+L antibody {ABflo® 594} (STJS001190)

STJS001190

GENERAL INFORMATION


Product Type	Secondary antibodies
Short Description	
Applications	IF/ICC/FC
Host/Source	Goat
Reactivity	

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	ABflo® 594
Purification	Affinity purification
Dilution Range	IF/ICC:1:50-1:200 FC:1:100-1:500
Formulation	PBS with 0.02% Sodium Azide, 1% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	

TARGET INFORMATION


Gene ID	
Gene Symbol	
Uniprot ID	
Immunogen	
Immunogen Region	
Specificity	Mouse IgG
Immunogen Sequence	




Immunofluorescence analysis of NIH/3T3 cells using ABflo® 594-conjugated Goat Anti-Mouse IgG (H+L) antibody (STJS001190) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using ABflo® 594-conjugated Goat Anti-Mouse IgG (H+L) antibody (STJS001190) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using ABflo® 594-conjugated Goat Anti-Mouse IgG (H+L) antibody (STJS001190) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Flow cytometric analysis of Positive antibody (STJ11103697) (2.5 µg/mL) in various cells (orange) compare to Mouse isotype control (blue) and non-staining control (red). The secondary antibody used was ABflo® 594-conjugated Goat Anti-Mouse IgG (H+L) (STJS001190) at 1:100.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081