

Goat Anti-IgG H+L antibody {ABflo® 647} (STJS001211)
STJS001211

GENERAL INFORMATION


Product Type	Secondary antibodies
Short Description	
Applications	IF/ICC/FC
Host/Source	Goat
Reactivity	

PRODUCT PROPERTIES

Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	ABflo® 647
Purification	Affinity purification
Dilution Range	IF/ICC:1:100-1:800 FC:1:100-1:800
Formulation	PBS with 0.025% Sodium Azide, 0.75% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	


TARGET INFORMATION

Gene ID	
Gene Symbol	
Uniprot ID	
Immunogen	
Immunogen Region	
Specificity	Mouse IgG
Immunogen Sequence	



Confocal imaging of U-2 OS cells using Alpha-Tubulin Mouse mAb (STJ11107556, dilution 1:400) followed by a further incubation with ABflo® 647-conjugated Goat Anti-Mouse IgG (H+L) (STJS001211, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of U-2 OS cells using AA-Tubulin Mouse mAb (STJ11107556, dilution 1:400) followed by a further incubation with ABflo® 647-conjugated Goat Anti-Mouse IgG (H+L) (STJS001211, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Flow cytometric analysis of Positive antibody Human Calctonin R (2.5 Mu g/mL) in various cells (orange) compare to Rabbit isotype control (blue) and non-staining control (Red). The secondary antibody used was ABflo® 647-conjugated AffinPure Goat Anti-Mouse IgG (H+L) (STJS001211) at 1:100.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081