

## Anti-BFAR antibody (1-140) (STJ110606)

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Rabbit polyclonal antibody anti-BFAR (1-140) is suitable for use in Western Blot and Immunofluorescence.

Applications WB, IF Host/Source Rabbit

Reactivity Human, Mouse, Rat

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID Concentration

Conjugation Unconjugated Purification Affinity purification Dilution Range WB 1:500-1:2000

IF 1:50-1:100

Formulation PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.

Isotype IgG

Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 51283 Gene Symbol BFAR

Uniprot ID BFAR\_HUMAN


Immunogen Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of human BFAR (NP\_057645.1).

Immunogen Region 1-140 Specificity


Immunogen Sequence




Western blot analysis of extracts of various cell lines, using BFAR antibody (STJ110606) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 12:100 graph and 12:100 graph 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



Immunofluorescence analysis of C6 cells using BFAR Polyclonal Antibody (STJ110606) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using BFAR Polyclonal Antibody (STJ110606) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using BFAR Polyclonal Antibody (STJ110606) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.