

Anti-Phospho-TP53-S37 antibody (STJ11101144)

ST.I11101144

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Rabbit polyclonal antibody anti-Phospho-p53-S37 is suitable for use in Western Blot and Immunohistochemistry.

Applications WB, IHC Host/Source Rabbit

Reactivity Human, Mouse, Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration Und

Conjugation Unconjugated Affinity purification Dilution Range WB 1:500-1:2000

IHC 1:50-1:200

Formulation PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.

Isotype IgG

Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.


TARGET INFORMATION

Gene ID 7157
Gene Symbol TP53
Uniprot ID P53 HUMAN


Uniprot ID P53_HUMAN

Immunogen A phospho specific peptide corresponding to residues surrounding S37 of human p53.

Immunogen Region Specificity Immunogen Sequence



Western blot analysis of evtracts of various cell lines, using Phospho-p53-837 antibody (STJ11101144) at 11000 dilution. Both 293T cells and C5 cells were treated by Calyculin A (100 nM) at 37 °C for 30 minutes after serum-starvation overnight. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-p5-35-37 antibody (STJ11011144) at 1:100. dilution, NIH/3T3 cells were treated by Calyculin A (100 n/M) at 37 °C for 30 minutes after seum-starvation overnight. Secondary antibody: HIPP Goat Anti-rabbi [9G (H-L) at 1:10000 dilution. Lysates/profeins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1809.