

## Anti-IDH2 antibody [ARC0567] (STJ11101307) STJ11101307

## **GENERAL INFORMATION**

 Product Type
 Primary antibodies

 Short Description
 Rabbit monoclonal antibody anti-IDH2 is suitable for use in Western Blot, Immunohistochemistry and Immunofluorescence.

 Applications
 WB, IHC, IF

 Reactivity
 Human, Mouse, Rat

## **PRODUCT PROPERTIES**

 Clonality
 Monoclonal

 Clone ID
 ARC0567

 Concentration
 Unconjugated

 Purification
 Affnity purification

 Dilution Range
 WB 1:500-1:200

 IHC 1:50-1:200
 IF 1:50-1:200

 IF 3:50-1:200
 IF 3:50-1:200


 Isotype
 JBS containing 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH7.3.

 Isotype
 IgG


 Storage Instruction
 Store in a freezer at-20°C and avoid freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 3418 Gene Symbol IDH2 Uniprot ID IDHP\_HUMAN Immunogen Region Specificity Immunogen Sequence



Western blot analysis of extracts of various cell lines, using IDH2 rabbit monoclonal antibody (STU11101307) at 1:1000 dilution. Secondary antibody: HRP Goat Antirabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per Iane. Blocking buffer 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 3s.



tibody (STJ11101307) at dilution of 1:100 (40x lens). rform microwave antigen retrieval with 10 mM PBS iffer pH 7. 2 before commencing with munohistochemistry staining protocol.



Immunofluorescence analysis of C6 cells using IDH2 rabbit monoclonal antibody (STJ11101307) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Immunofluorescence analysis of NIH-3T3 cells using IDH2 rabbit monocional antibody (STJ11101307) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081