Anti-COPS5 antibody [ARC0895] (STJ11101993)

STJ11101993

GENERAL INFORMATION

Product Type Primary antibodies
Short Description Rabbit monoclonal antibody anti-JAB1 is suitable for use in Western Blot, Immunohistochemistry and Immunofluorescence.
Applications WB, IHC, IF
Host/Source Rabbit
Reactivity Human, Mouse, Rat

PRODUCT PROPERTIES

Clonality Monoclonal
Clone ID ARC0895
Concentration
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB 1:500-1:2000
IHC 1:50-1:200
IF 1:50-1:200
ChIP 1:50-1:200
Formulation PBS containing 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH7.3.
Isotype IgG
Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.

TARGET INFORMATION

Gene ID 10987
Gene Symbol COPS5
Uniprot ID CSN5_HUMAN
Immunogen A synthesized peptide derived from human JAB1/CSN5/COPS5
Immunogen Region
Specificity
Immunogen Sequence



Western blot analysis of extracts of various cell lines, using JAB1/CSN5/COPS5 rabbit monoclonal antibody (STJ11101993) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.


Immunohistochemistry of paraffin-embedded rat testis using JAB1/CSN5/COPS5 rabbit monoclonal antibody (STJ11101993) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.


Immunohistochemistry of paraffin-embedded human colon using JAB1/CSN5/COPS5 rabbit monoclonal antibody (STJ11101993) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.


Immunohistochemistry of paraffin-embedded mouse testis using JAB1/CSN5/COPS5 rabbit monoclonal antibody (STJ11101993) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081