

## Anti-ALDH1L1 antibody [ARC1432] (STJ11102338)

STJ11102338

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Rabbit monoclonal antibody anti-ALDH1L1 is suitable for use in Western Blot, Immunohistochemistry and

Immunofluorescence.

**Applications** WB, IHC, IF **Host/Source** Rabbit

Reactivity Human, Mouse, Rat

## **PRODUCT PROPERTIES**

Clonality Monoclonal Clone ID ARC1432

Concentration

Conjugation
Purification
Purification
Dilution Range
Unconjugated
Affinity purification
WB 1:500-1:2000

IHC 1:50-1:200 IF 1:50-1:200

Formulation PBS containing 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH7.3.

**Isotype** IgG


**Storage Instruction** Store in a freezer at-20°C and avoid freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 10840
Gene Symbol ALDH1L1
Uniprot ID AL1L1\_HUMAN

Immunogen A synthesized peptide derived from human ALDH1L1


Immunogen Immunogen Region Specificity Immunogen Sequence




Western blot analysis of extracts of various cell lines using ALDH1L1 rabbit monoclonal antibod (STJ11102338) at 1:1000 dilution. Secondary antibod (HRP Goat Anti-rabbit IgG (H+L) at 1:1000 dilutior Lysates/proteins: 25up per Jane. Blocking buffer: 39 nonfat dry milk in TBST. Detection: ECL Basic Kit Fxposure fire: 908.



Immunohistochemistry of paraffin-embedded rat live using ALDHL1 rabbit monoclonal authoric (STJ11102338) at dilution of 1:100 (40x lens). Perforor microwave antigen retrieval with 10 mM Tris/EDTX buffer pH 9. 0 before commencing wit immunohistochemistry staining protocol



Immunohistochemistry of paraffin-embedded huma liver cancer using ALDHL1 rabbit monoclonal antibod (STJ11102338) at dilution of 1:100 (40x lens). Perforor microwave antigen retrieval with 10 mM Tris/EDT buffer pH 9. 0 before commencing with immunohistochemistry statingn protocol.



Immunohistochemistry of paraffin-embedded mouss kidney using ALDH11 rabbit monoclonal antibody (STJ1110238) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9. 0 before commencing with immunohistochemistry stations contend to the property of the prope