

## Anti-Neuregulin-3 antibody (STJ11103119)

ST.J11103119

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** Rabbit monoclonal antibody anti-Neuregulin-3 (NRG3) is suitable for use in Western Blot.

Applications WB
Host/Source Rabbit

Reactivity Human, Mouse, Rat

## **PRODUCT PROPERTIES**

Clonality Monoclona

Clone ID

Concentration Conjugation Unconjugated Purification Affinity purification Dilution Range WB 1:500-1:2000

Formulation PBS containing 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH7.3.

Isotype IgG

Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 10718
Gene Symbol NRG3
Uniprot ID NRG3\_HUMAN

Immunogen A synthesized peptide derived from human Neuregulin-3 (Neuregulin-3 (NRG3))


Immunogen Region
Specificity

Immunogen Sequence

Specificity Immunogen Sequence



Western blot analysis of extracts of various cell lines using Neuregulin-3 (Neuregulin-3 (Neuregulin-3 (Neuregulin-3 (Neuregulin-3 (Neuregulin-3 (Neuregulin-3) extra (ST.J11103119) at 1:1000 dilution. Secondary antibody Hogo Coat Anti-rabbit IgG (H+L) at 1:1000 dilution Lysates/proteins: 25up per Jane. Blocking buffer: 39 nonfat dry milk in TBST. Detection: ECL Basic Kit Forosure fine: 60s.



Western blot analysis of extracts of various cell lines, using Neuregulin-3 (Neuregulin-3 (Nelargulin-3 (Nelargulin-3 (Nelargulin-3 (Nelargulin-3 (Nelargulin-3)) antibody; HT Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.