

Anti-NELFCD antibody (STJ11103577)

STJ111103577

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Rabbit polyclonal antibody anti-NELFCD is suitable for use in Western Blot and Immunohistochemistry.

Applications WB, IHC Host/Source Rabbit

Reactivity Human, Mouse, Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration

Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB 1:500-1:2000

IHC 1:50-1:200

Formulation PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.

Isotype IgG

Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.


TARGET INFORMATION

Gene ID 51497
Gene Symbol NELFCD
Uniprot ID NELFD_HUMAN


Immunogen Recombinant fusion protein containing a sequence corresponding to amino acids 355-599 of human NELFCD

(NP_945327.2).


Immunogen Region Specificity Immunogen Sequence




Western blot analysis of extracts of various cell lines using NELFCD antibody (STJ11103577) at 1:100 dilution. Secondary antibody: HRP Goat Anti-rabbit lgd (H+L) at 1:10000 dilution. Lysates/proteins: 25ug pe lane. Blocking buffer: 3% nonfat dry milk in TBST Detection: FCI Basic Kif. Exposure time: 10s.



immunonistochemistry of paramin-embedoded numai colon using NELFCD rabbit polyclonal antibod, (STJ11103577) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffe pH 6.0 before commencing with immunohistochemistry statistics protocol



Immunohistochemistry of paraffin-embedded mous spleen using NELFCD rabbit polyclonal antibod (STJ11103577) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffe pH 6. 0 before commencing with immunohistochemistr staining protocol.



Immunohistochemistry of paraffin-embedded rat testi using NELFCD rabbit polyclonal antibod (STJ11103577) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffe pH 6. 0 before commencing wit immunohistochemistry staining protocol.