Anti-DDDDK-Tag antibody (STJ11103696)

STJ11103696

GENERAL INFORMATION

Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-DDDDK-Tag is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and Immunoprecipitation.
Applications	WB, IHC, IF, IP
Host/Source	Rabbit
Reactivity	

PRODUCT PROPERTIES

Clonality	Polyclonal
Clone ID	
Concentration	
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB 1:1000-1:2000 IHC 1:50-1:100 IF 1:50-1:100 IP 1:50-1:100
Formulation	PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.
Isotype	IgG
Storage Instruction	Store in a freezer at-20°C and avoid freeze-thaw cycles.

TARGET INFORMATION

Gene ID	
Gene Symbol	
Uniprot ID	
Immunogen	A synthetic peptide corresponding to DDDDK tag.
Immunogen Region	
Specificity	
Immunogen Sequence	



Western blot analysis of 293T-pCMV3-GFP-Flag-Myc-HA-His protein using rabbit anti DDDDK-Tag polyclonal antibody (STJ11103696) at different dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of 293T cells transfected with Flag-c (mCherry) and 293T cells transfected with Flag-N (mCherry) use rabbit anti DDDDK-Tag polyclonal antibody (STJ11103696) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.

Immunofluorescence analysis of 293T cells transfected with Flag-N (mCherry) and untreated 293T cells use rabbit anti DDDDK-Tag polyclonal antibody (STJ11103696) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 200ug extracts of 293T cells using 3ug rabbit anti DDDDK-Tag antibody (STJ11103696). Western blot was performed from the immunoprecipitate using rabbit anti DDDDK-Tag antibody (STJ11103696) at a dilition of 1:1000.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081