

Anti-Myc-Tag antibody (C-Term) (STJ11103699)

STJ11103699

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-Myc-Tag (C-Term) is suitable for use in Western Blot, Immunofluorescence and Immunoprecipitation.
Applications	WB, IF, IP
Host/Source	Rabbit
Reactivity	

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB 1:1000-1:10000 IF 1:50-1:200 IP 1:50-1:200
Formulation	PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.
Isotype	IgG
Storage Instruction	Store in a freezer at -20°C and avoid freeze-thaw cycles.

TARGET INFORMATION


Gene ID	
Gene Symbol	
Uniprot ID	
Immunogen	A synthetic peptide corresponding to Myc tag.
Immunogen Region	C-Term
Specificity	
Immunogen Sequence	



Western blot analysis of extracts of normal 293T cells and 293T transfected with Myc-NLK protein, using Myc-Tag antibody (STJ11103699) at 1:5000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



Immunofluorescence analysis of HeLa-Myc-CandHeLa cells using rabbit anti Myc-Tag pAb-C-terminal (STJ11103699) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 300ug extracts of 293T cells using 3ug rabbit anti Myc-Tag pAb-C-terminal antibody (STJ11103699). Western blot was performed from the immunoprecipitate using rabbit anti Myc-Tag pAb-C-terminal antibody (STJ11103699) at a dilution of 1:5000.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081