

Anti-CDK2 antibody (STJ113540)

ST.1113540

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Rabbit polyclonal antibody anti-CDK2 is suitable for use in Western Blot, Immunohistochemistry and Immunoprecipitation.

Applications WB, IHC, IP Host/Source Rabbit

Reactivity Human, Mouse, Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration

Conjugation
Purification
Purification
Affinity purification
Dilution Range
WB 1:500-1:2000

IHC 1:50-1:100 IP 1:50-1:200

Formulation PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.

Isotype IgG


Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.

TARGET INFORMATION


Gene ID 1017 Gene Symbol CDK2 Uniprot ID CDK2_HUMAN

Immunogen A synthetic peptide of human CDK2


Immunogen Region Specificity Immunogen Sequence



Western blot analysis of extracts from normal (contra and CDK2 knockout (KO) 293T cells, using CDK antibody (STJ113540) at 1:1000 dilution. Seconda antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:1000 (dilution. Lysates/proteins: 25ug per lane. Blockin buffer: 3% nonfat dry milk in TBST. Detection: EC Basic Kit. Exposure time: 103.



Western blot analysis of extracts of HeLa cells, using DK2 antibody (STJ1113540) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H-L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: CCL Basic Kit. Exposure time: 10s.



Immunohistochemistry of paraffin-embedded human esophageal cancer using CDK2 rabbit polyclonal antibody (STJ113540) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 ml Tris/EDTA buffer pH 9, 0 before commencing with immunohistochemistry staining portocot.