

## Anti-EGFR antibody (1100-1200) (STJ113559)

STJ113559

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Rabbit polyclonal antibody anti-EGFR (1100-1200) is suitable for use in Western Blot.

Applications WB Host/Source Rabbit

Reactivity Human, Mouse, Rat

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

Concentration

Conjugation
Purification
Dilution Range

Unconjugated
Affinity purification
WB 1:500-1:1000

Formulation PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.

Isotype IgG

Storage Instruction Store in a freezer at-20°C and avoid freeze-thaw cycles.

## **TARGET INFORMATION**


Gene ID 1956
Gene Symbol EGFR

Uniprot ID EGFR\_HUMAN


Immunogen A synthetic peptide corresponding to a sequence within amino acids 1100-1200 of human EGFR (NP\_005219.2).

Immunogen Region 1100-1200


Specificity Immunogen Sequence



Mestern blot analysis of extracts from normal (control and EGFR knockout (KO) HeLa cells, using EGF antibody (STJ113559) at 1:3000 dilution. Secondal antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:1000 dilution. Lysates/proteins: 25ug per lane. Blockin buffer: 3% nonfat drv milk in TBST. Detection: EG



Western blot analysis of extracts of various cell lines using EGFR antibody (STJ113559) at 1:1000 dilution Secondary antibody: HRP Goat Anti-rabbit 1gG (H-L) a 1:10000 dilution. Lysates/proteins: 25ug per lane Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic KIt. Exposure time: 10s.



Western blot analysis of extracts of Rat brain, using EGFR antibody (STJ113559) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBS1. Detection: ECL Enhanced Kit. Exposure time: 1800.