

## Anti-IGF2BP2 antibody (330-599) (STJ116159)

STJ116159

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-IGF2BP2 (330-599) is suitable for use in Western Blot, Immunohistochemistry and Immunofluorescence.
<b>Applications</b>	WB, IHC, IF
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB 1:500-1:2000 IHC 1:50-1:100 IF 1:50-1:100
<b>Formulation</b>	PBS containing 0.02% Sodium Azide, 50% Glycerol, pH7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store in a freezer at -20°C and avoid freeze-thaw cycles.

### TARGET INFORMATION


<b>Gene ID</b>	<a href="#">10644</a>
<b>Gene Symbol</b>	<a href="#">IGF2BP2</a>
<b>Uniprot ID</b>	<a href="#">IF2B2_HUMAN</a>
<b>Immunogen</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 330-599 of human IGF2BP2/IMP2 (NP_006539.3).
<b>Immunogen Region</b>	330-599
<b>Specificity</b>	
<b>Immunogen Sequence</b>	



Western blot analysis of extracts of various cell lines, using IGF2BP2/IMP2 antibody (STJ116159) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



Immunofluorescence analysis of U-2 OS cells using IGF2BP2/IMP2 antibody (STJ116159) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of rat testis cells using IGF2BP2/IMP2 antibody (STJ116159) at dilution of 1:100. Blue: DAPI for nuclear staining.