

Anti-CUL5 antibody (N-Term) (STJ13100099)

STJ13100099

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Nz White Rabbit polyclonal antibody anti-Cullin 5 (N-Term) is suitable for use in Immunohistochemistry and Western Blot research applications.
Applications	IHC, WB
Host/Source	NZ White Rabbit
Reactivity	Human, Rat, Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	
Conjugation	Unconjugated
Purification	Whole serum
Dilution Range	A dilution of 1:300 to 1:2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications.
Formulation	Shipped as lyophilised. Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.
Isotype	IgG
Storage	Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term.
Instruction	When reconstituting, glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles.

TARGET INFORMATION


Gene ID	8065
Gene Symbol	CUL5
Uniprot ID	CUL5_HUMAN
Immunogen	A synthetic peptide from the n-terminal of human Cullin 5 (Vasopressin activated calcium-mobilizing receptor 1) conjugated to an immunogenic carrier protein was used as the antigen.
Immunogen Region	N-Term
Specificity	Specific for Cullin 5.
Immunogen Sequence	




IHC on rat spinal cord (paraffin section) using Rabbit antibody to n-terminal of Cullin 5 (STJ13100099) at 1:200 dilution incubated overnight at 4C, developed with DAB/Ni and counterstained with hematoxylin.



IHC on rat spinal cord (paraffin section) using Rabbit antibody to n-terminal of Cullin 5 (STJ13100099) at 1:200 dilution incubated overnight at 4C, developed with DAB/Ni and counterstained with hematoxylin.



IHC on rat spinal cord (paraffin section) using Rabbit antibody to n-terminal of Cullin 5 (Vasopressin activated calcium-mobilizing receptor 1) : whole serum (STJ13100099) at 1:200 dilution incubated overnight at 4C, developed with DAB/Ni and counterstained with hematoxylin.



IHC on rat spinal cord (paraffin section) using Rabbit antibody to n-terminal of Cullin 5 (Vasopressin activated calcium-mobilizing receptor 1) : whole serum (STJ13100099) at 1:200 dilution incubated overnight at 4C, developed with DAB/Ni and counterstained with hematoxylin.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
 St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081