

Anti-CDH2 antibody (850aa to C-Term) (STJ140028)

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-N-Cadherin (850aa to C-Term) is suitable for use in Western Blot and Immunofluorescence research

Description applications. Applications WB, IF Host/Source Goat

Reactivity Human, Rat, Mouse, Monkey, Canine

PRODUCT PROPERTIES

Clonality Polyclonal Clone ID Concentration 3 mg/mL

Conjugation Unconjugated

Purification This antibody is epitope-affinity purified from goat antiserum.

Dilution Range WB 1:500-1:2000

IF 1:25-1:250

Formulation PBS, 20% glycerol and 0.05% sodium azide.

Isotype IgG

Storage For continuous use, store at 2-8 C for one-two days. For extended storage, store in-20 C freezer. Working dilution samples should

Instruction be discarded if not used within 12 hours.

TARGET INFORMATION

Gene ID 1000 Gene Symbol CDH2

Uniprot ID CADH2_HUMAN

Immunogen Purified recombinant peptide derived from within residues 850 aa to C-terminus of human CDH2 produced in E. coli.


Immunogen 850aa to C-Term

Region


Specificity Detects endogenous levels of total N-cadherin protein by Western blot in the whole cell lysates (COS-7, HeLa, HUH and Hepa-6).

Immunogen

Sequence



Endogenous CDH2 detected with at 1:500 dilution lysates at 100 µg per lane and rabbit polyclonal to goat IgG (HRP) at 1:10000 dilution.



N-Cadherin antibody marking the intercalated disks (distal cell junctions in heart tissue green line in the middle) of Human iPS-derived cardiomycoytes Methanol fixation DAPI: DNA (nuclear) stain, SARC aAct: Sarcomeric a-Actinin, N-CAD: N-Cadherin This image is courtesy of Dr Zanella F (University of California)