

## Anti-FTCD antibody (50aa N-Term) (STJ140093)

STJ140093

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Goat polyclonal antibody anti-Formimidoyltransferase cyclodeaminase (50aa N-Term) is suitable for use in Western Blot and Immunofluorescence research applications.
<b>Applications</b>	WB, IF
<b>Host/Source</b>	Goat
<b>Reactivity</b>	Human, Rat, Mouse, Monkey, Canine

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	3 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	This antibody is epitope-affinity purified from goat antiserum.
<b>Dilution Range</b>	WB 1:500-1:2000 IF 1:25-1:250
<b>Formulation</b>	PBS, 20% glycerol and 0.05% sodium azide.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	For continuous use, store at 2-8 C for one-two days. For extended storage, store in -20 C freezer. Working dilution samples should be discarded if not used within 12 hours.

### TARGET INFORMATION


<b>Gene ID</b>	10841
<b>Gene Symbol</b>	FTCD
<b>Uniprot ID</b>	FTCD_HUMAN
<b>Immunogen</b>	Purified recombinant peptide derived from within residues 50 aa to the N-terminus of human FTCD produced in E. coli.
<b>Immunogen Region</b>	50aa N-Term
<b>Specificity</b>	Detects a band of 55-60 kDa by Western blot in the following canine, human, monkey, mouse, rat whole cell lysates.
<b>Immunogen Sequence</b>	




Immunofluorescence – anti-FTCD antibody at 1:100 dilution in NHI:3T3 cells were fixed with methanol and permeabilized with 0.1% saponin



Immunofluorescence – anti-FTCD antibody at 1:100 dilution in NHI:3T3 cells were fixed with methanol and permeabilized with 0.1% saponin



Immunofluorescence – anti-FTCD antibody at 1:100 dilution in NHI:3T3 cells were fixed with methanol and permeabilized with 0.1% saponin



Immunofluorescence – anti-FTCD antibody at 1:100 dilution in NHI:3T3 cells were fixed with methanol and permeabilized with 0.1% saponin

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081