

## Anti-BCL2 antibody (41-54 aa) [124] (STJ180017)

STJ180017

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Mouse monoclonal antibody anti-BCL2 (41-54 aa) is suitable for use in Immunohistochemistry research applications.
<b>Applications</b>	IHC-p
<b>Host/Source</b>	Mouse
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES

<b>Clonality</b>	Monoclonal
<b>Clone ID</b>	124
<b>Concentration</b>	
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purified
<b>Dilution Range</b>	1:100-200
<b>Formulation</b>	Buffer with protein carrier and preservative
<b>Isotype</b>	IgG1/Kappa
<b>Storage</b>	Store at 2-8°C for up to 24 months. Predilute: Ready to use, no reconstitution necessary. Concentrate: Use dilution range and appropriate lab-standardized diluent. Stability after dilution: 7 days at 24°C, 3 months at 2-8°C, 6months at -20°C.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	596
<b>Gene Symbol</b>	BCL2
<b>Uniprot ID</b>	BCL2_HUMAN
<b>Immunogen</b>	A synthetic peptide, aa 41-54 (GAAPAGIFSSQPG-Cys) of human bcl-2 protein
<b>Immunogen Region</b>	41-54 aa
<b>Specificity</b>	Positive control: Tonsil or lymph node
<b>Immunogen Sequence</b>	



Human reactive lymph node stained with anti-BCL2 antibody using peroxidase-conjugate and DAB chromogen. Note cytoplasmic and membrane staining of mantle B-cells and perfollicular T-cells, and no staining of reactive B-cells in the germinal centers.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081