

## Anti-AMACR+p63+HMW CK antibody [13H4+ZM70+34 Beta E12] (STJ180270)

STJ180270

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit, Mouse monoclonal antibody anti-AMACR+p63+HMW CK is suitable for use in Immunohistochemistry research applications.
<b>Applications</b>	IHC-p
<b>Host/Source</b>	Rabbit, Mouse
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES

<b>Clonality</b>	Monoclonal
<b>Clone ID</b>	13H4+ZM70+34 Beta E12
<b>Concentration</b>	
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purified
<b>Dilution Range</b>	1:50
<b>Formulation</b>	Buffer with protein carrier and preservative
<b>Isotype</b>	IgG, IgG1 + IgG2a
<b>Storage Instruction</b>	Store at 2-8°C for up to 24 months. Predilute: Ready to use, no reconstitution necessary. Concentrate: Use dilution range and appropriate lab-standardized diluent. Stability after dilution: 7 days at 24°C, 3 months at 2-8°C, 6months at -20°C.

### TARGET INFORMATION

<b>Gene ID</b>	23600
<b>Gene Symbol</b>	AMACR
<b>Uniprot ID</b>	AMACR_HUMAN
<b>Immunogen</b>	Human AMACR (P504S) , p63, and CKHMW (34 Beta E12) polypeptides
<b>Immunogen Region</b>	
<b>Specificity</b>	Positive control: Prostate intraepithelial neoplasia (PIN)
<b>Immunogen Sequence</b>	



Human prostate carcinoma stained anti-AMACR, anti-p63 and anti-HMW CK 34βE12 antibodies using peroxidase-conjugate and DAB chromogens. Note cytoplasmic and luminal circumferential staining of carcinoma glands with AMACR and basal cell staining with p63 and HMW CK 34βE12.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081