

## Anti-Flotillin 1 antibody (C-Term) (STJ70184)

STJ70184

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-Flotillin 1 (C-Term) is suitable for use in ELISA, Western Blot, Immunofluorescence and Flow

Cytometry research applications.

Applications Pep-ELISA, WB, IF, FC

Host/Source Goat

Reactivity Human, Dog, Pig, Cow

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

**Concentration** 0.5 mg/mL **Conjugation** Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

**Dilution Range** WB-0.1-0.3μg/ml

IHC-5µg/ml

ELISA-antibody detection limit dilution 1:32000.

Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Isotype** IgG

Storage Store at-20 on receipt and minimise freeze-thaw cycles.

Instruction

## **TARGET INFORMATION**

Gene ID 10211

Gene Symbol FLOT1
Uniprot ID FLOT1\_HUMAN

Immunogen

Immunogen C-Term

Region

0-161111


Specificity This antibody is expected to recognize both reported isoforms (NP\_005794.1; NP\_001305804.1).

Immunogen SISQVNHKPLRTA


Sequence

260kDa 150kDa 150kDa 75kDa 50kDa 37kDa 25kDa 20kDa

STJ70184 (0. 1µg/ml) staining of K562 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



S1070164 (apg/mi) stammig of paramin embeddet Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6. HRP-staining.



STJ70184 Negative Control showing staining of paraffir embedded Human Kidney, with no primary antibody.