

Anti-IRAK4 antibody (N-Term) (STJ70645)

STJ70645

GENERAL INFORMATION


| | |
|--------------------------|---|
| Product Type | Primary antibodies |
| Short Description | Goat polyclonal antibody anti-IRAK4 (N-Term) is suitable for use in ELISA and Immunohistochemistry research applications. |
| Applications | Pep-ELISA, IHC |
| Host/Source | Goat |
| Reactivity | Human, Mouse |

PRODUCT PROPERTIES


| | |
|----------------------------|--|
| Clonality | Polyclonal |
| Clone ID | |
| Concentration | 0.5 mg/mL |
| Conjugation | Unconjugated |
| Purification | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Dilution Range | IHC-3.75µg/ml IF-Strong expression of the protein seen in the cytoplasm and microtubules of HeLa cells. 10µg/ml ELISA-antibody detection limit dilution 1:32000. |
| Formulation | 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. |
| Isotype | IgG |
| Storage Instruction | Store at-20 on receipt and minimise freeze-thaw cycles. |

TARGET INFORMATION


| | |
|---------------------------|--------------|
| Gene ID | 51135 |
| Gene Symbol | IRAK4 |
| Uniprot ID | IRAK4_HUMAN |
| Immunogen | |
| Immunogen Region | N-Term |
| Specificity | |
| Immunogen Sequence | NKPITPSTYVRC |



STJ70645 (1µg/ml) staining of HeLa (A) and Jurkat (B) cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



STJ70645 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml), showing cytoplasmic and microtubule staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml).



STJ70645 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (1µg/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081