

Anti-Nucleophosmin/NPM1 antibody (C-Term) (STJ70677)

STJ70677

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-Nucleophosmin/NPM1 (C-Term) is suitable for use in ELISA and Western Blot research

applications.

Applications Pep-ELISA, WB

Host/Source Goat

Reactivity Human, Mouse, Rat, Dog, Cow

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration 0.5 mg/mL Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the

immunizing peptide.

Dilution Range IHC-5µg/ml

IF-Strong expression of the protein seen in the nucleoli of HeLa cells. $10\mu\text{g/ml}$

ELISA-antibody detection limit dilution 1:8000.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin.}$

Isotype la

Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION

Gene ID 4869 Gene Symbol NPM1

Uniprot ID NPM_HUMAN


Immunogen

Immunogen Region C-Term


Specificity

Immunogen QEAIQDLWQWRKSL


Sequence



STJ70677 (1µg/ml) staining of NIH3T3 (A) , (0. 1ug/ml) Daudi (B) , Jurkat (C) and (0. 01ug/ml) K562 (D) cel lysate (35µg protein in RIPA buffer). Detected by chemilumipascence



STJ70677 (1µg/ml) staining of Mouse Spleen lysate 35µg protein in RIPA buffer). Detected by hemiluminescence.



STJ70677 Immunofluorescence analysis of paraformatchyde fixed Hela cells, permeabilized with 0. 15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nucleoil staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat Ifoug/ml) followed by Alexa Fluor 488 secondary (10ug/ml) followed by Alexa Fluor 488 secondary