

## Anti-MUC13 antibody (Internal) (STJ71483) STJ71483

## **GENERAL INFORMATION**

Host/Source Goat


Product Type Primary antibodies Short Description Goat polyclonal antibody anti-MUC13 (Internal) is suitable for use in ELISA and Western Blot research applications. Applications Pep-ELISA, WB Reactivity Human

## **PRODUCT PROPERTIES**


Clonality Polyclonal Clone ID Concentration 0.5 mg/mL Conjugation Unconjugated Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Dilution Range IF-Strong expression of the protein seen in the cytoplasm of HepG2 and Caco-2 cells. 10µg/ml ELISA-antibody detection limit dilution 1:4000. Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Isotype IgG Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 56667 Gene Symbol MUC13 Uniprot ID MUC13\_HUMAN Immunogen Immunogen Region Internal Specificity Immunogen DCKSDLQRPNPQ Sequence



noflucrescence analysis of ixed HepG2 cells, permeabilized Primary incubation 1hr (10ug/ml) Fluor 488 secondary antibody cytoplasmic staining. The nuclear plasmic staining. The ni egative control: Unimmu llowed by Alexa Fluor nized



Immunofluorescence analy nyde fixed Caco-2 cells, perm Triton. Primary incubation 1hr Alexa Fluor 488 secondary owing cytoplasmic staining. Th (blue). Negative control: Unin Oug/mi) followed by Alexa F naide 15% d by ) , st (10ug/ml) antibody nimr Flu

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081