

Anti-OAZ antibody (N-Term) (STJ71544)

STJ71544

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-OAZ (N-Term) is suitable for use in ELISA and Western Blot research applications.

Applications Pep-ELISA, WB Host/Source Goat

Reactivity Human, Mouse

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration 0.5 mg/mL **Conjugation** Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the

immunizing peptide.

Dilution Range IF-Strong expression of the protein seen in the nuclei of A431 cells. 10µg/ml

FC-Flow cytometric analysis of Kelly cells. 10ug/ml ELISA-antibody detection limit dilution 1:2000.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin.}$

Isotype IgG

Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION


Gene ID 23090 Gene Symbol ZNF423 Uniprot ID ZN423_HUMAN

Immunogen
Immunogen Region N-Term


Specificity

Immunogen EPECDQKTSRALEDR

Sequence



STJ/1544 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0. 15% Triton. Primary incubation 1hr (10uy/ml) followed by Alexa Fluor 488 secondary antibod, (2ug/ml), showing nuclear staining. Actin filaments were stained with phaliolidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) = followed by Alexa Fluor 488 secondary.



STJ71544 Flow cytometric analysis of paraformaldehyde fixed Kelly cells (blue line) permeabilized with 0.5% firton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml), IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary artibody.