

Anti-Doublecortin-Mouse antibody (Internal) (STJ72841)

STJ72841

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-Doublecortin-Mouse (Internal) is suitable for use in ELISA and Western Blot research applications.

Description

Applications Pep-ELISA, WB Host/Source Goat

Reactivity Human, Mouse, Rat

PRODUCT PROPERTIES

Clonality Polyclonal Clone ID

Concentration 0.5 mg/mL
Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

Dilution WB-0.5-2µg/ml

Range IF-Strong expression of the protein seen in the cytoplasm of HepG2 cells. 5µg/ml

ELISA-antibody detection limit dilution 1:128000.

Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA

Isotype IgG

Storage Store at-20 on receipt and minimise freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 1641 Gene Symbol DCX

Uniprot ID DCX_HUMAN

Immunogen

Immunogen Internal


Regio

Specificity This antibody is expected to recognize all reported mouse isoforms (NP_001103692.1; NP_001103694.1; NP_034155.2). Reported


variants represent identical protein: NP_001103692.1, NP_001103693.1

Immunogen PKASPTPQKTSAKS

Sequence



STJ72841 (0. 5µg/ml) staining of Mouse fetal Brain lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



STJ/2841 Immunofluorescence analysis o paraformalcelyvde fixed HepG2 cells, permeabilized with 0. 15% Triton. Primary incubation 1hr (Suyām followed by Alexa Fluor 488 secondary antibod (1ug/ml), showing cytoplasmic staining. The nucleas stain is DAPI (blue). Negative control: Unimmunized goat IgG (11ug/ml) followed by Alexa Fluor 488 spaces.