

Anti-MYD88, Biotinylated antibody (Internal) {Biotin} (STJ73268)

GENERAL INFORMATION

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-MYD88, Biotinylated (Internal) is suitable for use in ELISA and Western Blot research

applications

Applications Pep-ELISA, WB Host/Source Goat Reactivity Human

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration 0.5 mg/mL Conjugation Biotin

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the

immunizing peptide.

Dilution Range WB-0.1-0.3µg/ml

ELISA-antibody detection limit dilution 1:16000. Formulation 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA

Isotype IgG

Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION

Gene ID 4615

Gene Symbol MYD88

Uniprot ID MYD88_HUMAN . Immunogen

Immunogen Region Internal

Specificity This antibody is expected to recognize reported isoforms 1, 2 and 3 (NP_001166038.1; NP_002459.2; NP_001166039.1).

Immunogen IKYKAMKKEFP.

Sequence

250kDa 150kDa 100kDa 75kDa

50kDa 37kDa

25kDa 20kDa

ed STJ70667 (0. 2µg/ml) staining lysate (35µg protein in RIPA buffer its parental non-biotinylated produ n was 1 hour. Detectinescence, using streptavidin-HRP