

## Anti-TRPM2 antibody (Internal) (STJ73598)

STJ73598

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-TRPM2 (Internal) is suitable for use in ELISA and Western Blot research applications.

Applications Pep-ELISA, WB

Host/Source Goat

Reactivity Human, Mouse, Cow

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

**Concentration** 0.5 mg/mL **Conjugation** Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the

immunizing peptide.

Dilution Range IHC-3.75µg/ml

IF-Strong expression of the protein seen in the nuclear membranes and cytoplasm of A431 cells.  $10\mu g/ml$ 

ELISA-antibody detection limit dilution 1:16000.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$ 

**Isotype** IgG

Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 7226 Gene Symbol TRPM2


Uniprot ID TRPM2\_HUMAN

Immunogen
Immunogen Region Internal


Specificity

Immunogen QQKQRPEQKIED

Sequence



STJ73598 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with formal control of the paraformal control unirumunized goat IgG (10ug/mi) followed by Alexa Fluor 486 secondary artibody (2ug/mi).



STJ73598 Flow cytometric analysis of paraformaldehyde fixed Jurkat cells (blue line) , permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG black line) followed by Alexa Fluor 488 secondary