

## Anti-CTGF antibody (111-123) (STJ73609)

ST.173609

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Description Goat polyclonal antibody anti-CTGF (111-123) is suitable for use in ELISA and Western Blot research applications.

Applications Pep-ELISA, WB

Host/Source Goat

Reactivity Human, Mouse, Rat, Dog, Pig, Cow

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

Concentration 0.5 mg/mL Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the

immunizing peptide.

Dilution Range IHC-5µg/ml

IF-Strong expression of the protein seen in the cytoplasm of HepG2 and NIH3T3 cells.  $10\mu\text{g/ml}$ 

ELISA-antibody detection limit dilution 1:16000.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$ 

**Isotype** IgG

Storage Instruction Store at-20 on receipt and minimise freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 1490 Gene Symbol CCN2

Uniprot ID CCN2\_HUMAN


Immunogen

Immunogen Region 111-123


Specificity

Immunogen RSGESFQSSCKYQ


Sequence



STJ78699 Immunofluorescence analysis o paraformalcelyde fixed HepG2 cells, permeabilized with 0. 15% firtion. Primary incubation 1hr (10ug/ml followed by Alexa Fluor 488 secondary antibod (2ug/ml), showing cytoplasmic staining. The nuclea stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibod (2ug/ml).



STJ78609 Immunofluorescence analysis of oprarformalcelyyde fraxe MIHST3 cells, permeabilized with 0. 15% friton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary artibody (2ug/ml).



STJ73609 (5µg/ml) staining of paraffin embedded Human Thyroid. Steamed antigen retrieval with citrate