

## Anti-Phospho-LAT-Tyr200 antibody (140-220) (STJ90487)

STJ90487

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Phospho-Linker For Activation Of T-Cells Family Member 1-Tyr200 (140-220) is suitable for use in Western Blot and ELISA research applications.
<b>Applications</b>	WB, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution</b>	WB 1:500-1:2000
<b>Range</b>	ELISA 1:10000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	<a href="#">27040</a>
<b>Gene Symbol</b>	<a href="#">LAT</a>
<b>Uniprot ID</b>	<a href="#">LAT_HUMAN</a>
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human LAT around the phosphorylation site of Tyr171 at amino acid range 171-220
<b>Immunogen Region</b>	140-220
<b>Specificity</b>	Phospho-LAT-Tyr200 polyclonal antibody (Linker For Activation Of T-Cells Family Member 1) binds to endogenous Linker For Activation Of T-Cells Family Member 1 at the amino acid region 140-220 only when phosphorylated at Tyr200.
<b>Immunogen Sequence</b>	



Western blot analysis of lysates from Jurkat cells treated with UV 15', using LAT (Phospho-Tyr171) Antibody. The lane on the right is blocked with the phospho peptide.