

Anti-ATF7 antibody (10-90 Internal) (STJ91755)

STJ91755

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-Cyclic Amp-Dependent Transcription Factor Atf-7 (10-90 Internal) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry and ELISA research applications.
Applications	WB, IHC-P, IF, ICC, ELISA
Host/Source	Rabbit
Reactivity	Human, Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	1 mg/mL
Conjugation	Unconjugated
Purification	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
Dilution Range	WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:20000
Formulation	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	11016
Gene Symbol	ATF7
Uniprot ID	ATF7_HUMAN
Immunogen	The antiserum was produced against synthesized peptide derived from human ATF7 at amino acid range 41-90
Immunogen Region	10-90 Internal
Specificity	ATF7 polyclonal antibody (Cyclic Amp-Dependent Transcription Factor Atf-7) binds to endogenous Cyclic Amp-Dependent Transcription Factor Atf-7 at the amino acid region 10-90 Internal.
Immunogen Sequence	



Western blot analysis of lysates from HeLa and HUVEC cells, using ATF7 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of various cells using ATF-7 Polyclonal Antibody

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081