

## Anti-CST2 antibody (30-110 Internal) (STJ92619)

STJ92619

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Cystatin-Sa (30-110 Internal) is suitable for use in Immunohistochemistry, Immunofluorescence and ELISA research applications.
<b>Applications</b>	IHC-P, IF-P, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Rat, Mouse

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution Range</b>	IHC 1:100-1:300 ELISA 1:40000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	1470
<b>Gene Symbol</b>	CST2
<b>Uniprot ID</b>	CYTT_HUMAN
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CST2 at amino acid range 11-60
<b>Immunogen Region</b>	30-110 Internal
<b>Specificity</b>	CST2 polyclonal antibody (Cystatin-Sa) binds to endogenous Cystatin-Sa at the amino acid region 30-110 Internal.
<b>Immunogen Sequence</b>	



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using CST2 Antibody. The picture on the right is blocked with the synthesized peptide.