

Anti-FSHR antibody (180-260 Internal) (STJ93160)

STJ93160

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-Follicle-Stimulating Hormone Receptor (180-260 Internal) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and ELISA research applications.
Applications	WB, IHC-P, IF-P, ELISA
Host/Source	Rabbit
Reactivity	Human, Mouse, Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	1 mg/mL
Conjugation	Unconjugated
Purification	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
Dilution Range	IHC 100-300 WB 1:500-1:2000 ELISA 1:5000
Formulation	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	2492
Gene Symbol	FSHR
Uniprot ID	FSHR_HUMAN
Immunogen	The antiserum was produced against synthesized peptide derived from human FSHR at amino acid range 211-260
Immunogen Region	180-260 Internal
Specificity	FSHR polyclonal antibody (Follicle-Stimulating Hormone Receptor) binds to endogenous Follicle-Stimulating Hormone Receptor at the amino acid region 180-260 Internal.
Immunogen Sequence	



Western blot analysis of lysates from Jurkat and HUVEC cells, using FSHR Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:100 (4°C overnight), 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval, 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:100 (4°C overnight), 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval, 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

Immunohistochemical analysis of paraffin-embedded Human Amygdala. 1, Antibody was diluted at 1:100 (4°C overnight), 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval, 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081